



CREaTE

Canterbury Research and Theses Environment

Canterbury Christ Church University's repository of research outputs

<http://create.canterbury.ac.uk>

Please cite this publication as follows:

Coulton, S., Clift, S. M., Skingley, A. and Rodriguez, J. (2015) Effectiveness and cost-effectiveness of community singing on the mental health related quality of life of the older population: a randomized controlled trial. *British Journal of Psychiatry*, 207 (3). pp. 250-255. ISSN 0007-1250.

Link to official URL (if available):

<http://dx.doi.org/10.1192/bjp.bp.113.129908>

This version is made available in accordance with publishers' policies. All material made available by CReaTE is protected by intellectual property law, including copyright law. Any use made of the contents should comply with the relevant law.

Contact: create.library@canterbury.ac.uk



Effectiveness and cost-effectiveness of community singing on the mental health related quality of life of the older population: A randomized controlled trial.

Simon Coulton¹, Stephen Clift², Ann Skingley², John Rodriguez³.

¹ Centre for Health Service Studies, University of Kent, Canterbury, UK, CT2 7NZ. ² Sidney De Haan Research Centre for Arts and Health, Canterbury Christ Church University, North Holmes Road, Canterbury, UK, CT1 1QU. ³ NHS Kent and Medway, Templar House, Tannery Lane, Ashford, UK, TN23 1PL.

Corresponding author: Simon Coulton, Centre for Health Service Studies, University of Kent, Canterbury, UK, CT2 7NZ. s.coulton@kent.ac.uk

Abstract

Background: As the population ages, older people account for a greater proportion of the health and social care budget. While some research has been conducted on the use of music therapy for specific clinical populations, little rigorous research has been conducted looking at the value of community singing on the mental health-related quality of life of older people.

Aims: To evaluate the effectiveness and cost-effectiveness of community group singing for a population of older people in England.

Method: A pilot pragmatic individual randomized controlled trial comparing group singing versus usual activities in those aged 60 years or more.

Results: 258 participants were recruited across 5 centres in East Kent. At 6 months post-randomisation significant differences were observed in terms of mental health related quality of life measured using the SF12, mean difference 2.35 (95% CI 0.06 to 4.76) in favour of group singing. In addition the intervention was found to be marginally more cost-effective than usual activities. At 3 months significant differences were observed in terms of mental health components of quality of life (4.77; 2.53 to 7.01) anxiety (-1.78; -2.5; -1.06) and depression (-1.52; -2.13 to -0.92).

Conclusions: Community group singing appears to have a significant effect on mental health related quality of life, anxiety and depression and may be a useful intervention to maintain and enhance the mental health of the older population.

Declaration of Interest: Potential conflict of interest is listed at the end of the manuscript.

Trial Registration ISRCTN 62404401

INTRODUCTION

The increasing number of older people in the United Kingdom (UK) account for a significant proportion of health and social care service use(1) and this requires novel approaches to maintain and promote the mental and physical health of this population. There is evidence that maintaining an active lifestyle mentally, physically and socially is important in contributing to 'successful ageing', wellbeing, and the ability to remain living independently(2). Previous evidence based reviews of interventions aimed at maintaining or enhancing mental health and health related quality of life in older people found little evidence for a variety of group based interventions including exercise, tai chi and reminiscence groups(3) but a systematic review by Bridle et al(4) highlighted the benefits of tailored exercise in reducing depression in older people. Recent years have witnessed a growing recognition of the value of participatory arts activities in improving the mental health and overall wellbeing of older people(5). Within the area of music, previous research with older people focussed upon the impact of music listening(6, 7) or music therapy(8), rather than music as part of everyday experience(9). Clift et al(10) conducted a systematic mapping of non-clinical research studies focusing specifically on participatory singing and found only two studies using standardized measures within controlled trials for older people(5, 11). Both studies identified improvements in mental health for participants in singing groups, but have serious methodological limitations in terms of lack of justification for sample size and failure to randomize to intervention or control group, issues that limit the value of the evidence. Further reviews have pointed to the value of singing as a therapeutic intervention for older people

with long-term physical health conditions(12, 13), including small pilot randomised controlled trials of singing lessons for people with Chronic Obstructive Pulmonary Disease in clinical settings(14, 15).

The focus of the current study is the evaluation of an innovative community singing initiative, the 'Silver Song Club Project', which provides opportunities for older people to come together to sing with the support of professional musicians. Some 40 such clubs currently exist, mainly in South East England, which are managed by a third sector organization, Sing For Your Life Ltd (SFYL). Initially we completed a qualitative, process-orientated evaluation which suggested potential positive benefits across psychological, cognitive, social, and physical domains(16) and this provided a basis upon which to develop a randomised controlled trial to evaluate the effectiveness and cost-effectiveness of community singing on the mental and physical health related quality of life of older people.

Aims of the study

1. To assess the effectiveness for older people of active engagement in community singing on measures of mental and physical health related quality of life, depression and anxiety.
2. To evaluate the cost-effectiveness for older people of active engagement in community singing.

Hypotheses

Primary hypothesis, stated as null hypothesis:

Singing groups for older people are no more effective than usual activities in increasing mental health-related quality of life in older people measured six months after randomisation assessed using the York SF12.

Secondary hypotheses, stated as null hypotheses:

1. Singing groups for older people are no better at reducing anxiety and depression when compared with usual activities at six months after randomisation assessed using the Hospital and Anxiety depression Scale.
2. Singing groups for older people are no more effective than usual activities in increasing physical health-related quality of life in older people assessed six months after randomisation using the York SF12
3. Singing groups for older people are no more cost-effective than usual activities.

METHODS

A prospective pilot pragmatic randomised controlled trial in which eligible, consenting participants were randomised with equal probability to either singing group participation or usual activities. Randomisation was conducted by a secure remote randomisation service independent of the research team. Randomisation employed random permuted blocks of variable length and was stratified by centre and gender. Participants were followed up at 3 and 6 months by post. The study was approved by the Surrey NHS ethics committee (ref: 10/H1109/5) and registered (ISRCTN 62404401). The study was undertaken in accordance with the Declaration of Helsinki.

Sample size

The primary outcome measure was the mental health component of quality of life measured using the SF12 at 6 months post-randomisation. A clinically important difference on this dimension is estimated as a difference of 5 points between intervention and control group, equivalent to a medium effect size difference of 0.5(17). To detect this difference using a two-tailed test, alpha of 0.05 and power at 80% requires 63 participants in each of the two arms, a total of 126. We anticipated 5 singing groups and 5 controls and needed to take account of any clustering effect in calculating sample size. We used a conservative estimate of intra-class correlation coefficient of 0.02, similar to populations other community dwelling older populations and a harmonic mean cluster size of 12. This inflated the required sample size by a factor of 1.2, a total of 154. Previous research suggested the loss to follow up at 6 months for this population would be in the order of 20% and this further inflated the sample size to 184, 92 in each of the intervention and control groups.

Participants

As the intervention focussed on maintaining or enhancing mental health status experiencing current mental health issues was not a specific inclusion criteria. The study was publicised widely within the local areas. Researchers attended day centres and other venues where older people met for group activities to provide information on the study. In addition advertisements were placed in the local media, general practices and community venues. Inclusion and exclusion criteria were kept to a minimum to maximise the generalizability

of the population. All those expressing an interest, aged 60 years or more were eligible to participate. Only those unable to provide informed consent were excluded.

Procedure

All individuals indicating an interest in taking part in the study were sent a baseline questionnaire, an information sheet outlining the purpose of the study and a consent form to be returned with the questionnaire. Eligible and consenting participants were randomised to either a singing group or usual activities.

Intervention

Control group

Individuals in the control group continued with their normal activities. In order to address any potential resentful demoralisation members of the control group were informed that they would be welcome to join a singing group at the end of the research study, after completion of the primary outcome assessment at 6 months.

Intervention group

The Silver Song Club model is an established format for participative singing for older people that was selected as one of three UK examples of good practice selected for inclusion for the Health Pro Elderly international project(18). Details are available on the SFYL website

(http://www.singforyourlife.org.uk/sites/default/files_new/SSC%20FormativevEvalu

[lation%20Summary.pdf](#)). Trained and experienced facilitators under the guidance of SFYL met to compile a fourteen week 90 minute programme comprising songs from different eras and a variety of genres. This was followed by a series of 'unification' meetings, to ensure that all facilitators were aware of how to access the material and deliver it in the same way (e.g. accompaniment, musical key, acquiring copyright). The programme was developmental, progressing from singing melody lines to harmonising, layering and singing in rounds. Chime bars were also introduced where appropriate and there was an opportunity for participants to request particular songs. All clubs delivered the same programme concurrently and at the end of the fourteen weeks the clubs disbanded. A programme manager, who made unannounced visits to each group during the intervention period, monitored fidelity. A songbook was produced for the trial and a register of attendees was maintained.

Study measures

Primary outcome

The primary outcome measure was mental health-related quality of life assessed using the York SF-12(19) at 6 months post-randomisation. The SF-12 contains 12-items addressing both mental and physical health components of quality of life and has established psychometric properties including reliability, validity and sensitivity to change.

Secondary outcome measures

The SF-12 was also used to generate physical health-related components of quality of life. Anxiety and depression was measured using the Hospital

Anxiety and Depression Scale(20). This scale is validated for community samples and provides both an increasing severity score ranging from 1 to 21, with higher scores indicating greater severity.

Economic outcome measures

Health utility was measured using the EQ-5D(21). This is a short, 3-level, 5-dimensional instrument allows the generation of Quality Adjusted Life Years. It is routinely used in the economic evaluation of health care and recommended for cost-effectiveness analyses. Health and social care service utilisation was measured using a specially designed service use postal questionnaire used previously in a number of evaluations including older populations(22, 23) and measures units of health and social care resources including general practice visits, social care involvement, inpatient stays and outpatient attendance.

All outcomes were measured at baseline, prior to randomisation and then at 3 and 6 months by post. If a participant failed to respond to a follow-up questionnaire a reminder and additional questionnaire was sent 4 weeks after the scheduled follow-up date. In addition we collected process measures consisting of individual attendance at singing groups and fidelity information on the delivery of singing groups.

Analysis

As a study of effectiveness the primary analysis was by intention-to-treat where participants were analysed as part of their allocated group irrespective of the actual treatment received. The primary outcome measure, SF-12

mental components at 6 months was analysed using an analysis of covariance adjusting for baseline age and gender which are known covariates, as the intervention involved groups we adjusted the analysis using the Huber-White Sandwich estimation technique to generate robust standard errors. Secondary outcomes were analysed in a similar manner.

The incremental cost effectiveness of singing groups compared with usual activities were assessed from a health and social care perspective in accordance with National Institute of Clinical Excellence guidelines(24). The costs associated with setting-up and running singing groups were assessed from the actual local costs including the cost associated with premises and managerial overheads. Units of service utilisation in the 6 months prior to, and 6 months after randomisation were assessed from the service use questionnaire and the net costs for each arm of the study were derived by multiplying these by national sources of unit costs(25), as all costs were collected within a 12-month period no discounting was applied. The EQ5D was used with population values to calculate the quality adjusted life year (QALY) change using the area under the curve method(26). As economic data tends to be skewed, we used an established bootstrapping technique, resampling with replacement, to derive more robust confidence interval estimates(27). We divided the differences in the net costs for each arm by the difference in QALY gains to yield an incremental cost-effectiveness ratio. We estimated the sampling distribution from 1000 bootstrapped samples and derived cost-effectiveness acceptability curves(28). These curves plot the

resulting probability that one arm is better than the other against the maximum policy makers may be willing to pay for an additional QALY.

RESULTS

Sample characteristics

Recruitment to the study took place across 5 localities in East Kent. A total of 393 potential participants expressed an interest and were sent information on the study and a baseline questionnaire. Of these 258 (66%) were eligible and consented to participate in the study and 127 (49%) were allocated to the control and 131 (51%) allocated to the intervention. Follow-up rates at 3 and 6 months were 222 (86%) and 204 (79%) respectively and no differential follow-up rate between the groups was observed. Of those allocated to the singing groups 106 (81%) attended at least 50% of the sessions and attendance was similar across all centres. A full CONSORT diagram is provided in figure 1.

[Insert figure 1]

Baseline demographics and outcome measures are provided in table 1. The mean age was 69 (SD 7.14); the majority were female (84%) and white (98%). No statistical differences in baseline demographics or baseline outcome measures were observed between the groups.

[Insert table 1]

Primary outcome.

At baseline SF12 mental health related quality of life was similar across the groups 50.0 (95% CI 47.9 to 52.2) and 48.8 (46.8 to 50.8) for the control and

intervention groups respectively. While these were similar in the control group at 6 months, 49.9 (48.2 to 51.7) they had improved in the intervention group, 52.3 (50.7 to 54.0). The mean difference between intervention and control at 6 months was 2.35 (0.06 to 4.76) and this was significant ($p=0.05$) (table 2).

[Insert table 2]

Secondary outcomes

At 6 months no significant differences were observed between the groups in terms of SF12 physical components of health-related quality of life, anxiety or depression [table 2]. At three months significant differences between the groups were observed in terms of; mental components of SF12 health-related quality of life, mean difference intervention compared to control 4.77 (2.53 to 7.01), anxiety -1.78 (-2.50 to -1.06) and depression -1.52 (-2.13 to -0.92). No other significant differences were observed between the groups at 3 months (table 2).

Economic outcomes

The cost of implementing and training staff to conduct the singing groups are shown in table 3. Training costs were estimated over a 12-month period where the average facilitator would deliver 80 sessions, 2 per week, in order to avoid an over-estimation of training costs. The total cost per session was estimated at £176.84 and the cost per participant over 14 sessions estimated at £18.88.

[Insert table 3]

Service use was measured at baseline and 6 months and costs of units consumed derived from national sources. Service use costs were estimated for those followed-up only and are presented in table 4. Service use costs increased in both groups between baseline and 6 months but while the increase was greater in the intervention group, £315.89 versus £281.14 for the control group, this difference was not significant. Participants in the control group gained 0.008 QALY's between baseline and 6 months compared with a gain of 0.023 QALY's in the intervention group, the difference between the groups of 0.015 (95% CI 0.014 to 0.016) was significant.

[Insert table 4]

A cost-effectiveness acceptability curve was constructed (figure 2). This indicated that at a willingness to pay (WTP) threshold of zero the control group would be the preferred economic option. At a WTP threshold of £20 000 the intervention has a 60% probability of being the more cost-effective option and at recommended WTP thresholds of £30 000(24) this probability increases to 64%.

[Insert figure 2]

DISCUSSION

The reported study is the first pragmatic randomised controlled trial of community singing groups for older people focussing on their mental health and quality of life. The interest shown in the groups and the willingness of participants to engage in singing groups is a clear indication of both feasibility and acceptability of community singing for the older populations. Our primary

hypothesis explored the potential benefits of singing groups on maintaining the mental health related quality of life of older people. The results suggest that participation in singing groups confer significant benefit in terms of mental aspects of quality of life derived using SF12 and appear cost-effective when compared to usual activities six months after randomisation and 3 months after the groups had ceased to meet. No differences were observed at 6 months in terms of physical aspects of quality of life, anxiety or depression. At 3 months, at the end of the intervention levels of anxiety and depression were significantly lower in the singing group. This suggests that the greatest benefit occurs while participants are engaged in singing groups and continued access to singing groups may confer important benefits on the mental health of the older population.

Qualitative feedback from participants through written comments and interviews was highly positive. People indicated their enjoyment of the experience and highlighted the benefits on mental health, wellbeing and social relationships.

A clear marker of the value participants placed on the singing groups is the fact that four of the five groups established for research purposes were reinstated at the end of the six month follow-up and continue to meet and have grown in membership with support from a new charity Living Lively (<http://www.livinglively.org.uk/>). The study adds weight to the notion that meaningful, social and pleasurable activities can confer mental health benefits to participants identified in other studies of music therapy(8).

In conclusion the provision of opportunities to meet and sing together provide an opportunity to maintain and enhance the mental health of older people that is cost-effective and acceptable to the population, and should be considered as an important element in any public mental health strategy for this population.

Limitations of our study include the fact that it was conducted in one geographical area where the population is predominantly white British. We do not know, therefore, whether our findings could be generalized to other areas with different demographic characteristics. The groups also ran for a relatively short period of time and it is possible that longer involvement in singing could lead to more substantial and sustained benefit. While the study reported was pragmatic we did not explore in detail the processes of change that may underpin any observed changes and understanding these processes may be important in understanding the relationship between group musical activity and improvements in wellbeing.

It may be the case that any group activity confers similar benefits and that singing groups are just one form of group activity and further research is needed to address the relative effects of group singing versus other group activities. Yet it is important to note the ease with which the sample was recruited and the high levels of engagement would suggest that group singing is both feasible and acceptable to the older population. In addition the design of the study involved a waiting list control group, where participants allocated

to the control group had the intervention made available at the end of the study. The reasons for this involved addressing an issue of resentful demoralisation. It may have been the case that the control group was perceived as a delayed intervention for some participants and this may have impacted on their reporting of the outcomes. If this was the case then the reported effects at 6 months may have been under-estimates of the true effect. In addition it is important to be clear that the sample was a self-selecting population, people who wanted to engage in singing groups, but the ease of recruitment and the numbers of expressions of interest suggest there are large numbers of individuals who would like to engage in singing groups if they were more widely available.

Further, as our focus was on mental health related quality of life the study population was not specifically experiencing severe mental health issues. We have undertaken an observational study of group singing for people with enduring and severe mental health issues over a year, and this demonstrated clinically important improvement in mental health outcomes. The next step in building on the pilot trial reported here should be a larger-scale multi-centre trial running over a longer period of time.

Acknowledgement

The study was funded by the National Institute for Health Research under the Research for Patient Benefit Programme. The views expressed herein do not necessarily reflect the funder. We wish to thank our collaborators and all participants who contributed to the conduct of this study.

Authors' contribution

All authors contributed to the design and development of the study protocol. Conduct of the project and interpretation of the study results. Authors have commented on successive drafts and all authors read and approved the final manuscript.

Conflict of interest

SC, AS, JR has no conflict of interest to declare. SCI is a board member of Sing For Your Life Ltd a not for profit third sector organisation who played a role in developing and implementing the intervention reported.

REFERENCES

1. Imison C. Future trends. Kings Fund, 2012.
2. Health Development Agency. Taking local action: improving the health and wellbeing of people in mid-life and beyond. NICE, 2004.
3. Van Malderen L, Mets T, Gorus E. Interventions to enhance the Quality of Life of older people in residential long-term care: a systematic review. *Ageing research reviews*. 2013; 12(1): 141-50.
4. Bridle C, Spanjers K, Patel S, Atherton NM, Lamb SE. Effect of exercise on depression severity in older people: systematic review and meta-analysis of randomised controlled trials. *The British journal of psychiatry : the journal of mental science*. 2012; 201(3): 180-5.
5. Cohen GD, Perlstein S, Chapline J, Kelly J, Firth KM, Simmens S. The impact of professionally conducted cultural programs on the physical health, mental health, and social functioning of older adults. *The Gerontologist*. 2006; 46(6): 726-34.
6. Coffman D. Music and quality of life in older people. *Psychomusicology*. 2006; 46(6): 76-88.
7. Stacey R, Brittain K, Kerr S. Singing for health: an exploration of the issues. *Health Educ*. 2002; 199(2): 132-29.
8. Erkkila J, Punkanen M, Fachner J, Ala-Ruona E, Pontio I, Tervaniemi M, et al. Individual music therapy for depression: randomised controlled trial. *The British journal of psychiatry : the journal of mental science*. 2011; 199(2): 132-9.
9. Sixsmith A, Gibson G. Music and the wellbeing of people with dementia. *Ageing and Soc*. 2006; 27(): 127 - 45.
10. Clift S, Nicols J, Raisbeck M, Whitmore C, Morrison I. Group singing, wellbeing and health: a systematic mapping of the research evidence. *UNESCO Jnl*. 2010; 2(1).
11. Houston D, McKee J, Carroll L, Marsh H. Using humour to promote psychological wellbeing in residential homes for older people. *Ageing and Mental Health*. 1998; 2(4): 328 - 32.
12. Clark I, Harding K. Psychosocial outcomes of active singing as a therapeutic intervention: a review of the literature. *Nordic Jnl of Mus Ther*. 2012; 21(1): 80 - 98.
13. Gick M. Singing, health and wellbeing: a health psychologist's review. *Psychomusicology*. 2011; 21(1): 176 - 207.
14. Lord VM, Cave P, Hume VJ, Flude EJ, Evans A, Kelly JL, et al. Singing teaching as a therapy for chronic respiratory disease--a randomised controlled trial and qualitative evaluation. *BMC pulmonary medicine*. 2010; 10: 41.
15. Lord VM, Hume VJ, Kelly JL, Cave P, Silver J, Waldman M, et al. Singing classes for chronic obstructive pulmonary disease: a randomized controlled trial. *BMC pulmonary medicine*. 2012; 12: 69.
16. Skingley A, Bungay H. The Silver Song Club Project: singing to promote the health of older people. *British journal of community nursing*. 2010; 15(3): 135-40.
17. Qalymetric. SF12 v2 Health survey users manual. Qalymetric, 2010.
18. Billings J, Brown P. HealthProElderly - Evidence based guidelines for older people: United Kingdom national evaluation report. University of Kent, 2008.

19. Iglesias CP, Birks YF, Torgerson DJ. Improving the measurement of quality of life in older people: the York SF-12. *QJM : monthly journal of the Association of Physicians*. 2001; 94(12): 695-8.
20. Snaith R, Sigmond A. The hospital anxiety and depression scale. NFER, 1994.
21. EuroQoL Group. EuroQoL-a new facility for the measurement of health-related quality of life. *Health Pol*. 1990; 16(3): 199-208.
22. Richardson G, Bloor K, Williams J, Russell I, Durai D, Cheung WY, et al. Cost effectiveness of nurse delivered endoscopy: findings from randomised multi-institution nurse endoscopy trial (MINuET). *Bmj*. 2009; 338: b270.
23. Watson JM, Crosby H, Dale VM, Tober G, Wu Q, Lang J, et al. AESOPS: a randomised controlled trial of the clinical effectiveness and cost-effectiveness of opportunistic screening and stepped care interventions for older hazardous alcohol users in primary care. *Health technology assessment*. 2013; 17(25): 1-158.
24. NIHSCE. Guide to the methods of technology appraisal. NIHSCE, 2013.
25. Curtis L. The unit costs of health and social care. (ed PSSRU). University of Kent, 2007.
26. Brooks R, Rabin R, de Charro F. The measurement and valuation of health status using EQ-5D: a European perspective. Kluwer Academic Publishers, 2003.
27. Efron B, Tibshirami R. An introduction to the bootstrap. Chapman & Hall, 1993.
28. Fenwick E, Claxton K, Sculpher M. Representing Uncertainty: the role of the cost-effectiveness acceptability curves. *Health Econ*. 2001; 10: 779-87.

Table 1: Baseline description of the sample

	Overall (n=258)	Control (n=127)	Intervention (n=131)
Demographics			
Mean age (SD)	69.2 (7.14)	69.5 (7.13)	69.2 (7.18)
Female n (%)	214 (83.9)	108 (87.1)	106 (80.9)
Smoking n (%)	11 (4.3)	3 (2.4)	8 (6.2)
White n (%)	250 (98.0)	120 (96.8)	130 (99.2)
Employed n (%)	25 (11.0)	9 (8.1)	16 (13.8)
Education after 16 n (%)	162 (62.8)	79 (64.8)	83 (63.8)
Outcome measures			
Mean SF12 – Physical score (SD)	39.4 (6.63)	39.8 (6.69)	39.1 (6.58)
Mean SF12 – Mental score (SD)	49.4 (11.7)	50.0 (11.9)	48.8 (11.5)
Mean EQ5D score (SD)	0.74 (0.22)	0.74 (0.22)	0.74 (0.22)
Mean HADS – Anxiety (SD)	6.40 (4.46)	6.41 (4.57)	6.40 (4.46)
Anxiety case n (%)	49 (19.1)	24 (19.0)	25 (19.1)
Mean HADS – Depression (SD)	4.62 (3.52)	4.28 (3.52)	4.95 (3.52)
Depression case n (%)	20 (7.8)	8 (6.3)	12 (9.2)

Table 2: Baseline, 3 and 6-month outcomes adjusted for baseline values, age and gender.

	Baseline	Month 3			Month 6		
	Mean (95% CI)	Mean (95% CI)	Mean Difference (95% CI)	p-value	Mean (95% CI)	Mean Difference (95% CI)	p-value
SF12 – Physical							
Control	39.8 (38.6; 40.9)	39.2 (38.3; 40.0)			39.6 (38.6; 40.7)		
Intervention	39.1 (37.9; 40.3)	40.0 (39.1; 40.8)	0.83 (-0.39; 2.05)	0.18	39.9 (38.7; 40.9)	0.26 (-1.75; 1.23)	0.73
SF12 – Mental							
Control	50.0 (47.9; 52.2)	50.7 (49.1; 52.3)			49.9 (48.2; 51.7)		
Intervention	48.8 (46.8; 50.8)	55.5 (53.9; 57.1)	4.77 (2.53; 7.01)	<0.01	52.3 (50.7; 54.0)	2.35 (0.06; 4.76)	0.05
HADS - Anxiety							
Control	6.41 (5.62; 7.20)	6.01 (5.41; 6.42)			5.83 (5.30; 6.36)		
Intervention	6.40 (5.62; 7.18)	4.14 (3.64; 4.64)	-1.78 (-2.50; -1.06)	<0.01	5.26 (4.75; 5.76)	-0.57 (-1.31; 0.16)	0.13
HADS - Depression							
Control	4.28 (3.67; 4.89)	4.15 (3.72; 4.56)			4.22 (3.71; 4.73)		
Intervention	4.95 (4.53; 5.57)	2.63 (2.21; 3.05)	-1.52 (-2.13; -0.92)	<0.01	3.69 (3.20; 4.18)	-0.53 (-1.24; 0.18)	0.14
EQ5D - QALY							
Control	0.76 (0.72; 0.81)	0.78 (0.74; 0.82)			0.77 (0.72; 0.82)		
Intervention	0.76 (0.71; 0.80)	0.80 (0.76; 0.85)	0.02 (0.01; 0.03)	0.05	0.78 (0.73; 0.83)	0.01 (0.01; 0.02)	0.01

Table 3: Implementation and training costs associated with singing groups

Resource	Unit Cost per session £	Cost per participant (n=131)
Training costs		
Facilitators (5 facilitators, 3 days at £75/day)	2.81 ¹	0.30
Facilitator expenses (5 facilitators, 3 journeys at £35/journey)	1.31 ¹²	0.14
Trainer (3 days at £100/day)	0.75 ³	0.08
Trainer expenses (3 journeys at £35/journey)	0.26 ⁴	0.03
Venue Hire (3 days @ £100/day)	0.75 ⁵	0.08
Capital Expenditure		
Hand Chimes	1.25 ⁶	0.13
Keyboard	1.75 ⁷	0.19
Song sheets	0.22 ⁸	0.02
Indirect cost		
Advertising	2.14 ⁹	0.23
Management	9.64 ¹⁰	1.03
Administration	12.86 ¹¹	1.37
Session costs		
Facilitator	75.00 ¹²	8.01
Facilitator expenses	35.00 ¹²	3.74
Venue Hire	30.00 ¹²	3.21
Refreshments	3.00 ¹²	0.32
Total	176.84	18.88

¹ Total training cost for facilitators is £1125. Training estimated per annum at 80 sessions delivered per facilitator, cost per session £2.81.

² Total travel cost for facilitators is £525. Training estimated per annum at 80 sessions delivered per facilitator, cost per session £1.31.

³ Total trainer cost estimated at £300 to train 5 facilitators. Training estimated per annum at 80 sessions delivered per facilitator, cost per session £0.75.

4. Total trainer travel estimated at £105 to train 5 facilitators. Training estimated per annum at 80 sessions delivered per facilitator, cost per session £0.26.
5. Venue hire estimated at £300. Training estimated per annum at 80 sessions delivered per facilitator, cost per session £0.75.
6. Hand chimes unit cost of £500. Expected use 5 year, £100 per year, expected utilization estimated at 80 session per annum, cost per session £1.25.
7. Keyboard unit cost of £700. Expected use 5 years, £140 per year, expected utilization estimated at 80 sessions per year, cost per session £1.75.
8. Song sheets unit cost of £180 across 5 groups, £36 per group. Expected use 2 years, £18 per year, expected utilization at 80 sessions per year, cost per session £0.22.
9. Advertising cost per group per year at £120. Cost over 3 months £30 per group, for 14 sessions estimated at £2.14 per session.
10. Management cost per group per year £540. Cost over 3 months £135 per group, for 14 sessions estimated £9.64 per session.
11. Administration cost per group per year £720. Cost over 3 months £180 per group, for 14 sessions estimated £12.86 per session.
12. Actual cost.

Table 4: Mean (SE) service use costs for the 6 months pre-baseline and 6 months post-baseline

	Social Care (£)	Primary Care (£)	Secondary Care (£)	Total (£)
Baseline				
Control	2.58 (1.57)	66.38 (7.83)	273.62 (64.70)	342.59 (67.39)
Intervention	4.06 (2.23)	60.45 (5.51)	229.58 (50.39)	294.09 (52.87)
Month 6				
Control	5.04 (3.05)	85.21 (8.66)	533.48 (126.91)	623.73 (131.16)
Intervention	3.24 (1.82)	78.16 (8.25)	528.58 (208.70)	609.98 (210.15)

Figure 1: Trial Consort Statement




Figure 2: Cost-effectiveness acceptability curve comparing the probability of cost effectiveness for intervention and control at different QALY valuations

